

# Skew Dyck paths having no peaks at level 1 (Sequence A128723)

Helmut Prodinger  
Stellenbosch University  
Department of Mathematical Sciences  
7602 Stellenbosch  
South Africa,  
and  
NITheCS  
(National Institute for Theoretical and Computational Sciences)  
South Africa  
[hproding@sun.ac.za](mailto:hproding@sun.ac.za)

## Abstract

Skew Dyck paths are a variation of Dyck paths, where additionally to steps  $(1, 1)$  and  $(1, -1)$  a south-west step  $(-1, -1)$  is also allowed, provided that the path does not intersect itself. Replacing the south-west step by a red south-east step, we end up with decorated Dyck paths. Sequence A128723 of the Encyclopedia of Integer Sequences considers such paths where peaks at level 1 are forbidden. We provide a thorough analysis of a more general scenario, namely partial decorated Dyck paths, ending on a prescribed level  $j$ , both from left-to-right and from right-to-left (decorated Dyck paths are not symmetric). The approach is completely based on generating functions.

## 1 Introduction

Skew Dyck paths are a variation of Dyck paths, where additionally to steps  $(1, 1)$  and  $(1, -1)$  a south-west step  $(-1, -1)$  is also allowed, provided that the path does not intersect itself. Replacing the south-west step by a red south-east step, we end up with decorated Dyck paths. Our earlier publication [2] studied such paths using generating functions.

Sequence A128723 considers such paths where peaks at level 1 are forbidden. These paths are the main objects of the present paper.




Figure 1: All 10 skew Dyck paths of length 6 (consisting of 6 steps).




Figure 2: The 10 paths redrawn, with red south-east edges instead of south-west edges.

The Figures 1, 2, 3 describe such paths of length 6.

We catch the essence of a decorated Dyck path using a state-diagram (Fig. 4):

It has three types of states, with  $j$  ranging from 0 to infinity; in the drawing, only  $j = 0..8$  is shown. The first layer of states refers to an up-step leading to a state, the second layer refers to a black down-step leading to a state and the third layer refers to a red down-step leading to a state.

If the dashed edge is present, the graph models skew (decorated) Dyck paths. Any path from the origin to a node on level  $j$  represents such a decorated Dyck path ending on level  $j$ . In particular, if  $j = 0$ , the path comes back to the  $x$ -axis. Note that the syntactic rules of forbidden patterns  $\nearrow$  and  $\searrow$  can be clearly seen from the picture.

However, if the dashed edge is *not* present, it means that peaks at level 1 cannot be modeled by this graph, and that is what we want in the present paper.

We will work out generating functions describing all paths leading to a particular state. We will use the notations  $f_j, g_j, h_j$  for the three respective layers, from top to bottom. Although one could in principle compute all these functions separately, we are mainly interested in  $s_j = f_j + g_j + h_j$ , so we are interested in paths arriving on level  $j$  but we do not care in which way this final level has been reached. It is also clear that a path of length  $n$  leading to a state at level  $j$  must satisfy  $n \equiv j \pmod 2$ .

In a last section, the right-to-left model is briefly described. Then, red down-steps become blue up-steps.



Figure 3: The 6 paths without peaks on level 1.




Figure 4: Three layers of states according to the type of steps leading to them (up, down-black, down-red).

## 2 Generating functions and the kernel method

The functions depend on the variable  $z$  (marking the number of steps), but mostly we just write  $f_j$  instead of  $f_j(z)$ , etc.

The following recursions can be read off immediately from the diagram (Fig. 4):

$$\begin{aligned} f_0 &= 1, & f_{i+1} &= z f_i + z g_i, & i &\geq 0, \\ g_i &= z f_{i+1} + z g_{i+1} + z h_{i+1}, & i &\geq 1, \\ g_0 &= z g_{i+1} + z h_{i+1}, & i &\geq 1, \\ h_i &= z h_{i+1} + z g_{i+1}, & i &\geq 0. \end{aligned}$$

We can make a few direct observations:  $f_0 = 1$ ,  $f_1 = z + z g_0$ ,  $g_0 = h_0$ . The latter can be proved from combinatorial reasoning, since switching the last step from black to red resp. from red to black constitutes a bijection. This is a consequence of the fact that there are no peaks at level 1, otherwise the syntactic restrictions might be violated.

Now it is time to introduce *bivariate* generating functions:

$$F(z, u) = \sum_{i \geq 0} f_i(z) u^i, \quad G(z, u) = \sum_{i \geq 0} g_i(z) u^i, \quad H(z, u) = \sum_{i \geq 0} h_i(z) u^i.$$

Again, often we just write  $F(u)$  instead of  $F(z, u)$  and treat  $z$  as a ‘silent’ variable. Summing

the recursions leads to

$$\begin{aligned}\sum_{i \geq 0} u^i f_{i+1} &= \sum_{i \geq 0} u^i z f_i + \sum_{i \geq 0} u^i z g_i, \\ \sum_{i \geq 1} u^i g_i &= \sum_{i \geq 1} u^i z f_{i+1} + \sum_{i \geq 1} u^i z g_{i+1} + \sum_{i \geq 1} u^i z h_{i+1}, \\ \sum_{i \geq 0} u^i h_i &= \sum_{i \geq 0} u^i z h_{i+1} + \sum_{i \geq 0} u^i z g_{i+1}.\end{aligned}$$

This can be rewritten as

$$\begin{aligned}\frac{1}{u}(F(u) - 1) &= zF(u) + zG(u), \\ \sum_{i \geq 1} u^i g_i + g_0 &= \sum_{i \geq 1} u^i z f_{i+1} + \sum_{i \geq 1} u^i z g_{i+1} + z g_1 + \sum_{i \geq 1} u^i z h_{i+1} + z h_1, \\ \sum_{i \geq 0} u^i g_i &= \sum_{i \geq 1} u^i z f_{i+1} + \sum_{i \geq 0} u^i z g_{i+1} + \sum_{i \geq 0} u^i z h_{i+1}, \\ G(u) &= \frac{z}{u}[F(u) - f_0 - u f_1] + \frac{z}{u}[G(u) - g_0] + \frac{z}{u}[H(u) - h_0], \\ H(u) &= \frac{z}{u}(G(u) - G(0)) + \frac{z}{u}(H(u) - H(0)).\end{aligned}$$

Instead of working with 3 functions, we can reduce the system to just one equation (with the variable  $G$ ):

$$F = \frac{1 + zuG}{1 - zu}, \quad H = \frac{z(G - g_0 - h_0)}{u - z}.$$

Using this, we get

$$G = \frac{-z^3 u(u - z) + z(1 - zu)(2 + zu - z^2)g_0}{z(u - r_1)(u - r_2)}$$

with

$$r_1 = \frac{1 + z^2 + \sqrt{1 - 6z^2 + 5z^4}}{2z}, \quad r_2 = \frac{1 + z^2 - \sqrt{1 - 6z^2 + 5z^4}}{2z}.$$

Note that  $r_1 r_2 = 2 - z^2$ . Since the factor  $u - r_2$  in the denominator is “bad,” it must also cancel in the numerator. This is an essential step in the kernel method, see for instance our own survey [1]. This leads to the new version

$$G = \frac{-z^3(u - z + r_2) - z^2(-z^2 + zu + 1 + zr_2)g_0}{z(u - r_1)}.$$

Plugging in  $u = 0$  and solving the equation

$$G(z, 0) = g_0 = \frac{-z^3(-z + r_2) - z^2(-z^2 + 1 + zr_2)g_0}{z(-r_1)}$$

leads to

$$g_0 = \frac{1 - 2z^4 - 3z^2 - \sqrt{1 - 6z^2 + 5z^4}}{2(z^2 + 3)z^2}.$$

Knowing this, we know  $G$ , and thus  $F$  and  $H$ . As the first goal, we still set  $u = 0$ , thus considering paths coming back to the  $x$ -axis. Using Maple,

$$s_0 := f_0 + g_0 + h_0 = \frac{1 - z^4 - \sqrt{1 - 6z^2 + 5z^4}}{(z^2 + 3)z^2}.$$

## The conjecture

We write  $z^2 = x$ , since skew paths, as discussed here, must have an even number of steps. The function

$$y(x) = \frac{1 - x^2 - \sqrt{1 - 6x + 5x^2}}{x(x + 3)}$$

is the generating function of the sequence A128723:

1, 0, 2, 6, 22, 84, 334, 1368, 5734, 24480, 106086, 465462, 2063658, 9231084, 41610162, ...

GFUN, as described in [3], produces the algebraic equation that  $y(x)$  satisfies:

$$-(x - 1)(x - 2) + 3x + 2(1 - x^2)y - x(3 + x)y^2 = 0$$

and from this the differential equation

$$-(9x^2 + 5x^3 + 3 - 17x)xy' + (9x^2 + 7x - 5x^3 - 3)y + 3 + 9x^2 - 5x^3 - 7x$$

and finally from the differential equation the recursion for the coefficients  $s_n = [x^n]y(x)$ :

$$3(n + 4)s_{n+3} - (17n + 41)s_{n+2} + 9ns_{n+1} + 5(n + 1)s_n = 0.$$

An equivalent recursion was conjectured in the description of sequence A128723 [4].

## Partial paths

Another computation with Maple leads to

$$S(z, u) = F(z, u) + G(z, u) + H(z, u) = \frac{-z^4 - z^4g_0 - z^2g_0 + z^2 - 1}{z(u - r_1)}.$$

Further

$$\begin{aligned} s_j := [u^j]S(z, u) &= \frac{z^4 + z^4g_0 + z^2g_0 - z^2 + 1}{zr_1(1 - u/r_1)} \\ &= \frac{z^4 + z^4g_0 + z^2g_0 - z^2 + 1}{zr_1^{j+1}}. \end{aligned}$$

One sees the parity:  $j$  even/odd iff exponents are even/odd. If it is desired,  $1/r_1$  may be expressed by  $r_2$  (and a factor).

## Open-ended paths

We might allow *any* level as end-level of the path. In terms of generating functions, this means to consider  $S(z, 1)$ , viz.

$$S(z, 1) = \frac{-2z^5 - 3z^4 + z^3 - 5z^2 - 3z + 4 - (z^2 + 3z + 4)\sqrt{1 - 6z^2 + 5z^4}}{2z(3 + z^2)(z^2 + 2z - 1)}.$$

The sequence of coefficients

1, 1, 1, 2, 5, 8, 18, 31, 71, 126, 290, 527, 1218, 2253, 5223, 9796, 22763, 43170, 100502, 192347, ...

is not in the encyclopedia [4].

## 3 Reading the decorated paths from right to left

Since decorated paths are not symmetric, it makes sense to consider this scenario separately.



Figure 5: All 6 dual (=right-to-left) skew Dyck paths of length 6 (consisting of 6 steps), having no peak at level 1.

We catch the essence of a decorated (dual skew) Dyck path using a state-diagram:




Figure 6: Three layers of states according to the type of steps leading to them (down, up-black, up-blue).

Note that the syntactic rules of forbidden patterns  $\nearrow$  and  $\searrow$  can be clearly seen from the picture.

As in the earlier section, if the dashed edge is removed it means that the condition ‘no peak at level 1’ is modeled, which is what we need to do. Using the letters  $c_j, a_j, b_j$  (in

that order) for the generating functions to reach state  $j$  in the particular layer, we find the following recursions immediately from the diagram:

$$\begin{aligned} a_0 &= 1, & a_{i+1} &= za_i + zb_i + zc_i, & i &\geq 0, \\ b_0 &= zb_1, & b_i &= za_{i+1} + zb_{i+1}, & i &\geq 1, \\ c_{i+1} &= za_i + zc_i, & i &\geq 0. \end{aligned}$$

We introduce *bivariate* generating functions:

$$A(z, u) = \sum_{i \geq 0} a_i(z)u^i, \quad B(z, u) = \sum_{i \geq 0} b_i(z)u^i, \quad C(z, u) = \sum_{i \geq 0} c_i(z)u^i.$$

Summing the recursions leads to

$$\begin{aligned} A(u) &= \sum_{i \geq 0} u^i a_i = 1 + u \sum_{i \geq 0} u^i (za_i + zb_i + zc_i) \\ &= 1 + uzA(u) + uzB(u) + uzC(u), \\ \sum_{i \geq 0} u^i b_i &= \sum_{i \geq 1} u^i za_{i+1} + \sum_{i \geq 0} u^i zb_{i+1}, \\ B(u) &= \frac{z}{u} \sum_{i \geq 2} u^i a_i + \frac{z}{u} \sum_{i \geq 1} u^i b_i \\ &= \frac{z}{u} [A(u) - a_0 - ua_1] + \frac{z}{u} [B(u) - b_0], \\ \sum_{i \geq 1} u^i c_i &= uz \sum_{i \geq 0} u^i a_i + uz \sum_{i \geq 0} u^i c_i, \\ C(u) - c_0 &= uzA(u) + uzC(u). \end{aligned}$$

We have  $c_0 = 0$ ,  $a_0 = 1$ , and  $a_1 = z + zb_0$ . We may write

$$\begin{aligned} C(u) &= \frac{uzA(u)}{1 - uz}, \\ A(u) &= 1 + uzA(u) + uzB(u) + \frac{u^2 z^2 A(u)}{1 - uz} = \frac{1 + uzB(u)}{1 - uz - \frac{u^2 z^2}{1 - uz}} = \frac{1 - uz}{1 - 2uz} [1 + uzB(u)], \\ C(u) &= \frac{uz}{1 - 2uz} [1 + uzB(u)]. \end{aligned}$$

Solving for  $B(u)$ ,

$$B(u) = \frac{z(2u^2 z^2 + 2z^2 u^2 b_0 + b_0 z u - b_0)}{z(z^2 - 2)(u - r_1^{-1})(u - r_2^{-1})}.$$

We cancel the bad factor  $(u - r_1^{-1})$  out of the numerator:

$$B(u) = \frac{z(2r_1 uz + 2r_1 uz b_0 + b_0 r_1 + 2z + 2z b_0) r_2}{r_1 (z^2 - 2)(ur_2 - 1)}$$

Plugging in  $u = 0$  results in the equation

$$b_0 = \frac{-z(b_0 r_1 + 2z + 2z b_0) r_2}{r_1(z^2 - 2)}$$

and thus

$$b_0 = \frac{1 - z^4 - \sqrt{1 - 6z^2 + 5z^4}}{z^2(3 + z^2)} - 1,$$

as expected, since  $1 + b_0$  is the generating function of all skew Dyck paths without peaks at level 1.

Expressions for  $A(z, u) + B(z, u) + C(z, u)$  and  $[u^j](A(z, u) + B(z, u) + C(z, u))$  could be explicitly written, which we leave to the reader.

The open paths in this model are enumerated via

$$A(z, 1) + B(z, 1) + C(z, 1),$$

which is a long expression, with coefficients

$$1, 2, 4, 10, 24, 56, 134, 318, 764, 1824, 4390, 10520, 25346, 60878, 146768, \dots$$

which are again not in the OEIS [4].

Explicit formulæ for this model are a bit unpleasant, but easily regenerated using Maple.

## 4 Conclusion

In order to keep this paper short (and not boring) we refrained from working out many additional parameters. That might be a good project for graduate students.

## References

- [1] Helmut Prodinger. *The Kernel Method: A Collection of Examples*, Séminaire Lotharingien de Combinatoire, B50f (2004), 19 pp.
- [2] H. Prodinger, Partial skew Dyck paths—a kernel method approach, *arXiv:2108.09785* (2021).
- [3] B. Salvy and P. Zimmermann. GFUN: A Maple package for the manipulation of generating and holonomic functions in one variable. *ACM Trans. Math. Softw.*, 20(2): 163–177 1994.
- [4] Neil J. A. Sloane and The OEIS Foundation Inc. The on-line encyclopedia of integer sequences, 2022.



---

2010 *Mathematics Subject Classification*: Primary 05A15; Secondary 05A19.

*Keywords*: *Skew Dyck path, peak, forbidden pattern, generating function, kernel-method.*

---

(Concerned with sequence [A128723](#).)

---

Return to [Journal of Integer Sequences home page](#).